

## Western Marsh Harrier Preys on Herring Gull

Author(s): Gary R. Graves

Source: Journal of Raptor Research, 48(2):191-192. 2014.

Published By: The Raptor Research Foundation

DOI: <http://dx.doi.org/10.3356/JRR-13-00088.1>

URL: <http://www.bioone.org/doi/full/10.3356/JRR-13-00088.1>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

- COCKBURN, A. 2004. Mating systems and sexual conflict. Pages 81–101 in W. Koenig and J. Dickinson [EDS.], Ecology and evolution of cooperative breeding in birds. Cambridge University Press, Cambridge, U.K.
- CURSACH, J.A., C.G. SUAZO, R.P. SCHLATTER, AND J.R. RAU. 2012. Observaciones sobre el carancho negro *Phalcoeboenus australis* (Gmelin, 1788) en isla Gonzalo, Archipiélago Diego Ramírez, Chile. *Anales del Instituto de la Patagonia* 40:147–150.
- DWYER, J.F. AND S.G. COCKWELL. 2011. Social hierarchy of scavenging raptors on the Falkland Islands, Malvinas. *Journal of Raptor Research* 45:229–235.
- FERGUSON-LEES, J. AND D.A. CHRISTIE. 2001. Raptors of the world. Christopher Helm, London, U.K.
- KIMBALL, R.T., P.G. PARKER, AND J.C. BEDNARZ. 2003. Occurrence and evolution of cooperative breeding among the diurnal raptors (Accipitridae and Falconidae). *Auk* 120:717–729.
- KOENIG, W. AND J. DICKINSON. 2004. Ecology and evolution of cooperative breeding in birds. Cambridge University Press, Cambridge, U.K.
- MARÍN, M., A. KUSCH, D. OEHLER, AND S. DRIESCHMAN. 2006. Distribution, breeding and status of the Striated Caracara *Phalcoeboenus australis* (Gmelin, 1788) in southern Chile. *Anales del Instituto de la Patagonia* 34:65–74.
- MEIBURG, J.A. 2006. The biogeography of Striated Caracaras *Phalcoeboenus australis*. M.S. thesis, University of Texas at Austin, Austin, TX U.S.A.
- RAIMILLA, V., J. RAU, AND A. MUÑOZ-PEDREROS. 2012. Estado de arte del conocimiento de las aves rapaces de Chile: situación actual y proyecciones futuras. *Revista Chilena de Historia Natural* 85:469–480.
- REXER-HUBER, K. AND K.L. BILDSTEIN. 2013. Winter diet of Striated Caracara *Phalcoeboenus australis* (Aves, Polyborinae) at a farm settlement on the Falkland Islands. *Polar Biology* 36:437–443.
- SCHLATTER, R.P. AND G.M. RIVEROS. 1997. Historia natural del Archipiélago Diego Ramírez, Chile. *Serie Científica, Instituto Antártico Chileno* 47:87–112.
- SMITH, R.I.L. AND P.A. PRINCE. 1985. The natural history of Beauchêne Island. *Biological Journal of the Linnean Society* 24:233–283.
- STACEY, P.B. AND W.D. KOENIG. 1990. Cooperative breeding in birds: long-term studies of ecology and behavior. Cambridge University Press, Cambridge, U.K.
- STRANGE, I.J. 1996. The Striated Caracara *Phalcoeboenus australis* in the Falkland Islands. Philip Myers Press (Holding) Ltd., Warrington, U.K.
- TREJO, A. 2007. Identificación de especies y áreas prioritarias para el estudio de la reproducción de aves rapaces de Argentina. *Hornero* 22:85–96.
- VIRANI, M.Z. AND D.M. HARPER. 2004. A comparative study of the breeding behavior of the Augur Buzzard, *Buteo augur*, in two different land-use areas in southern Lake Naivasha, Kenya. *Ostrich* 75:11–19.
- WOODS, R.W. 2007. Distribution and abundance of the Striated Caracara *Phalcoeboenus australis* in the Falkland Islands–2006. Falklands Conservation, Stanley, Falkland Islands.

Received 10 September 2013; accepted 28 January 2014  
Associate Editor: Joan L. Morrison

*J. Raptor Res.* 48(2):191–192

© 2014 The Raptor Research Foundation, Inc.

## WESTERN MARSH HARRIER PREYS ON HERRING GULL

KEY WORDS: *Western Marsh Harrier*; *Circus aeruginosus*; *Herring Gull*; *Larus argentatus*; *predation*.

The Western Marsh Harrier (*Circus aeruginosus*) is one of the better known raptors in the western Palearctic (Cramp and Simmons 1980), and has been the subject of several diet studies (Schipper 1973, Underhill-Day 1985, Clarke et al. 1993, Brzeckiński and Żmihorski 2009, Cardador et al. 2012, Tornberg and Haapala 2013). Harriers feed on a wide range of small mammals and birds. Most avian prey is relatively small (10–675 g) but harriers occasionally take prey as large as 1000 g (e.g., Great Bittern [*Botaurus stellaris*]; Tornberg and Haapala 2013). However, prey items at the larger end of the mass spectrum tend to be

juvenile, sick, or disabled individuals (Schipper 1973). Nesting harriers in Finland frequently took Black-headed Gulls (*Chroicocephalus ridibundus*; 265 g) and less frequently Mew Gulls (*Larus canus*, 415 g; Tornberg and Haapala 2013). To my knowledge, there are no published records of Western Marsh Harrier preying on larger and more aggressive species of gulls.

On 8 September 2013, I watched a congregation of several thousand waterfowl, cormorants, shorebirds, Black-headed Gulls, Great Black-backed Gulls (*Larus marinus*) and Herring Gulls (*Larus argentatus*) from an observation

blind at Klydesøreservatet (55°34.8'N, 12°31.8'E) near Copenhagen, Denmark. Two Western Marsh Harriers made several passes over the mudflats and *Phragmites* beds before my attention was drawn to a commotion of Northern Lapwings (*Vanellus vanellus*) and Black-headed Gulls harassing a harrier that had captured a Herring Gull at the end of a narrow spit (12:24 H). I observed the action through a 20–60× scope from a distance of 205 m (measured with Google Earth Pro). The harrier had the gull's neck pinned to the mud with one foot, while the gull was on its back, feebly beating its wings, and slowly kicking its legs in a running fashion. The struggling gull failed to right itself and the harrier carried it to higher ground with the aid of a strong wind (approximately 28 km/hr). At the higher site, the harrier's feet and the gull's head were obscured by a dead thistle, so I could not see how the harrier finally dispatched the gull, which continued to kick until 12:33 H. The harrier began consuming bits of muscle soon afterwards and started plucking the gull's breast at 12:48 H. It fed until 12:52 H, stepped off the gull's carcass at 12:59 H, and preened nearby on the ground until I left the area at 13:09 H. Having missed the harrier's initial strike, I do not know whether the gull was acting normally beforehand or whether it showed signs of disease or incapacitation. However, the length of time (>9 min) required for the harrier to kill the gull suggested the gull was reasonably fit.

The harrier's golden crown and throat, and the absence of pale markings on the breast and leading edge of the wing indicated that it was a juvenile. The gull was in second- or third-year plumage, with dark mottling on the upper wing coverts. Herring Gulls from nominate populations in Norway vary in weight from 795 to 1440 g (Cramp and Simmons 1983), overlapping the size range of female harriers (540–1269 g) but not males (320–667 g; Cramp and Simmons 1980).

I thank two anonymous reviewers for helpful comments and the Smoketree Trust for support.—**Gary R. Graves** (email address: [gravesg@si.edu](mailto:gravesg@si.edu)), Department of Vertebrate Zoology, MRC-116, National Museum of Natural His-

tory, Smithsonian Institution, P.O. Box 37012, Washington, D.C. 20013 U.S.A, and Center for Macroecology, Evolution and Climate, University of Copenhagen, 2100 Copenhagen Ø, Denmark.

#### LITERATURE CITED

- BRZEKIŃSKI, M. AND M. ŻMIHORSKI. 2009. Nestling diet and parental provisioning behaviour in the Marsh Harrier (*Circus aeruginosus*). *Acta Zoologica Lituanica* 19:93–98.
- CARDADOR, L., E. PLANAS, A. VAREA, AND S. MAÑOSA. 2012. Feeding behaviour and diet composition of Marsh Harriers *Circus aeruginosus* in agricultural landscapes. *Bird Study* 59:228–235.
- CLARKE, R., A. BOURGONJE, AND H. CASTELIJNS. 1993. Food niches of sympatric Marsh Harriers *Circus aeruginosus* and Hen Harriers *C. cyaneus* on the Dutch coast in winter. *Ibis* 135:424–431.
- CRAMP, S. AND K.E.L. SIMMONS. [EDS.]. 1980. Handbook of the birds of Europe, the Middle East and North Africa: the birds of the western palearctic: hawks to bustards. Vol. 2. Oxford University Press, Oxford, U.K.
- AND ———. [EDS.]. 1983. Handbook of the birds of Europe, the Middle East and North Africa: the birds of the western palearctic: waders to gulls. Vol. 3. Oxford University Press, Oxford, U.K.
- SCHIPPER, W.J.A. 1973. A comparison of prey selection in sympatric harriers (*Circus* spp.) in western Europe. *Gerfaut* 63:17–120.
- TORNBERG, R. AND S. HAAPALA. 2013. The diet of the Marsh Harrier *Circus aeruginosus* breeding on the isle of Hailuoto compared to other raptors in northern Finland. *Ornis Fennica* 90:103–116.
- UNDERHILL-DAY, J.C. 1985. The food of breeding Marsh Harriers *Circus aeruginosus* in East Anglia. *Bird Study* 32:199–206.

Received 12 November 2013; accepted 10 January 2014

*J. Raptor Res.* 48(2):192–193

© 2014 The Raptor Research Foundation, Inc.

#### PREDATION BY GEOFFROY'S CAT ON FERRUGINOUS PYGMY-OWL IN CALDÉN SEMIARID FOREST, ARGENTINA

KEY WORDS: *Ferruginous Pygmy-Owl*; *Glaucidium brasilianum*; *Geoffroy's cat*; *Leopardus geoffroyi*; *Argentina*; *predation*.

The Ferruginous Pygmy-Owl (*Glaucidium brasilianum*) has a broad geographic distribution, occurring from the southwestern United States to Central America and South America (Holt et al. 1999, Proudfoot and Johnson 2000). It is

found in a variety of habitats from tropical and subtropical dry forest to semiarid open forest; it nests in natural cavities and is active diurnally (Proudfoot and Johnson 2000). Predation of Ferruginous Pygmy-Owl nestlings by raccoons