

Microbial diversity of extreme habitats in human homes

Amy M. Savage¹, Justin Hills², Katherine Driscoll³, Daniel J. Fergus⁴, Amy M. Grunden⁵ and Robert R. Dunn⁶

¹ Rutgers, The State University of New Jersey, Camden, United States

² Laboratory of Cellular and Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD, United States

³ Animal Management Department, The Wilds, Cumberland, OH, United States

⁴ Genomics and Microbiology, North Carolina Museum of Natural Sciences, Raleigh, NC, United States

⁵ Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC, United States

⁶ Department of Applied Ecology and Keck Center for Behavioral Biology, North Carolina State University, Raleigh, NC, United States

ABSTRACT

High-throughput sequencing techniques have opened up the world of microbial diversity to scientists, and a flurry of studies in the most remote and extreme habitats on earth have begun to elucidate the key roles of microbes in ecosystems with extreme conditions. These same environmental extremes can also be found closer to humans, even in our homes. Here, we used high-throughput sequencing techniques to assess bacterial and archaeal diversity in the extreme environments inside human homes (e.g., dishwashers, hot water heaters, washing machine bleach reservoirs, etc.). We focused on habitats in the home with extreme temperature, pH, and chemical environmental conditions. We found a lower diversity of microbes in these extreme home environments compared to less extreme habitats in the home. However, we were nonetheless able to detect sequences from a relatively diverse array of bacteria and archaea. Habitats with extreme temperatures alone appeared to be able to support a greater diversity of microbes than habitats with extreme pH or extreme chemical environments alone. Microbial diversity was lowest when habitats had both extreme temperature and one of these other extremes. In habitats with both extreme temperatures and extreme pH, taxa with known associations with extreme conditions dominated. Our findings highlight the importance of examining interactive effects of multiple environmental extremes on microbial communities. Inasmuch as taxa from extreme environments can be both beneficial and harmful to humans, our findings also suggest future work to understand both the threats and opportunities posed by the life in these habitats.

Subjects Biodiversity, Ecology, Microbiology

Keywords Community Ecology, Extreme environments, Human Homes, Interactive effects, Microbial diversity, Temperature, pH, Chemical

INTRODUCTION

The innovation of culture-independent, high-throughput sequencing techniques has facilitated the discovery of high microbial diversity in many habitats once considered inhospitable to life (*Rothschild & Mancinelli, 2001*). The species in these environments are frequent targets for the discovery of useful enzymes (*Niehaus et al., 1999; Van den Burg, 2003; Elleuche et al., 2014*), and studies of microbes living in extreme environments have

Submitted 16 March 2016
Accepted 29 July 2016
Published 13 September 2016

Corresponding author
Amy M. Savage,
amy.savage@rutgers.edu

Academic editor
Mauricio Rodriguez-Lanetty

Additional Information and
Declarations can be found on
page 13

DOI 10.7717/peerj.2376

© Copyright
2016 Savage et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

provided key insights into the evolution of microbial metabolism ([Valentine, 2007](#); [Hoehler & Jorgensen, 2013](#)). Often overlooked, however, is that the attributes that define many of the most extreme habitats on Earth, such as extremes of temperature, pH, water activity, or low nutrient levels, can also be found more immediate to everyday experience. Human homes, for example, contain microhabitats as hot, acidic, basic or salty as any encountered elsewhere on Earth ([Martin et al., 2015](#)).

We know of only two extreme habitats within homes where microbial diversity has been studied to date, and in both cases culture-dependent techniques were used. In 1973, Brock and Boylen discovered a species of the genus *Thermus* (*T. aquaticus*) living in hot water heaters. Species of this genus had previously been known only from hot springs ([Brock & Boylen, 1973](#)). In addition, studies have considered the biology of tap water. Tap water is hospitable in terms of its abiotic conditions (e.g., temperature, pH, toxicity) but is very low in nutrients and so was long assumed to be relatively devoid of life; until, that is, it was studied. Tap water has now been shown to contain many species of bacteria capable of surviving in low nutrient environments ([Kalmbach, Manz & Szewzyk, 1997](#); [Szewzyk et al., 2000](#); [Boe-Hansen et al., 2002](#)). If life exists in hot water heaters and tap water, it seems possible and even likely that many extreme habitats in homes sustain life. That the environmental extremes imposed by these conditions in homes (cold, hot, acidic, alkaline, wet or dry) delineate which species are present seems inevitable. That they are lifeless is unlikely.

Here, we used culture-independent, high-throughput sequencing to address the following questions: (1) What is the relative diversity of microbes (specifically, Bacteria and Archaea) under extremes of temperature, pH and chemical environments of southeast US homes and how does it compare to habitats without each extreme conditions? [Harrison et al. \(2013\)](#) recently argued that because many extreme environments include simultaneous extremes in multiple environmental factors, interactive effects of these multiple sources of extreme conditions are likely to be important determinants of microbial diversity in extreme environments. Therefore, we additionally asked (2) how do multiple, simultaneous extreme conditions influence microbial diversity in human homes? Finally, we asked (3) which bacterial and archaeal genera from the broader home ([Dunn et al., 2013](#)) fail to persist in extreme home habitats, and which microbial genera persist only in these extreme habitats?

METHODS

Sampling extreme home environments

We sampled extreme environments in six houses in the Raleigh-Durham metropolitan area ([Fig. S1](#)). In each house, we used dual-tipped sterile BBL™ CultureSwabs™ or 50 ml conical tubes to collect water from each of 10 standardized extreme locations in homes. The sites sampled in all six houses included environments that were extreme in terms of their temperature, pH and/or chemical environments ([Table S1](#)). Our assumptions concerning these sampling locations are based upon publicly available consumer resources regarding certain commercial and industrial requirements

(e.g., <http://www.nsf.org/consumer-resources/health-and-safety-tips/home-product-appliance-tips/sanitizing-dishwasher>, <http://energy.gov/energysaver/projects/savings-project-lower-water-heating-temperature>). For example, our sampling of dishwashers was influenced by the NSF/ANSI 184 standard for residential dishwashers to provide a final rinse at a temperature of at least 150 °F (65.6 °C). Additionally, temperature ranges for residential water heaters are 90–150 F (32–65.6 °C), depending on the manufacturer. Bleach receptacles in clothes washing machines would also be assumed to have a pH of 12 when bleach is present. Although the pH and chemical composition of laundry detergent and dishwasher detergent can be quite variable, manufacturing standards are generally within the 7–10 pH range. While measurements, opposed to assumptions, would be very useful, taking measurements of all the potential extreme axes under various sample sites in multiple homes was not feasible. All samples were preserved at –20 °C immediately after collection.

Isolating and identifying microbes in extreme home environments

Genomic DNA was extracted from all samples using the MoBio Power Soil DNA extraction kit (MoBio, Carlsbad, CA) as described previously (*Fierer et al., 2008*; *Lauber et al., 2009*). For swabs, the tips were placed in PowerBead tubes containing solution C1 and swirled vigorously for approximately 10 s to release contents and removed. Water samples were thawed and filtered using Corning 50 ml 0.22 µm cellulose acetate filters after which the filters were added to the PowerBead tubes. The extractions were subsequently performed as directed by the manufacturer, except that the final elution was performed in 50 µl of 70 °C C6 elution buffer. Because the water samples were frozen prior to filtering and extraction, the results reported for the water samples likely under-represents the true diversity of taxa in those environments.

We used methods described in *Bates et al. (2011)* to amplify bacterial and archaeal DNA from the samples collected from homes and six negative controls. Briefly, amplicons were produced by PCR with universal bacterial/archaeal 515F and 806R primers to which Roche 454 B pyrosequencing adapters had been added, as described in *Hulcr et al. (2012)*. The 515F primer contained an additional 12-bp barcode sequence for individual sample identification. All the samples were amplified by triplicate PCR reactions, cleaned using the UltraClean-htp 96-well PCR Clean-up kit (MoBio), and quantified with a Quant-iT PicoGreen dsDNA Assay kit (Invitrogen). Equimolar amounts of each sample were pooled into a single sample to sequence. DNA pyrosequencing was performed at Selah Clinical Genomics Center at Innovista (University of South Carolina, USA) using a Roche Genome Sequencer 454 FLX system to facilitate comparison to previous related work that utilized this platform (*Dunn et al., 2013*). Though these methods here do not distinguish living from recently dead cells, with the comparative approach used here, we presume that taxa frequently identified in one habitat but rare or absent in most others are likely surviving in the habitat from which they are frequently identified. The sequences were submitted to NCBI (SRA accession number [SRP071677](https://www.ncbi.nlm.nih.gov/sra/SRP071677)).

The QIIME analysis package (*Caporaso et al., 2010a*) was used to process and analyze the barcoded microbial (bacterial and archaeal) amplicon sequences. Sequences were

quality filtered to a minimum quality score of 25 with no unambiguous bases and sorted to each sample by the 12 bp barcodes. The 454 pyrosequencing produced 197,305 reads that passed the quality screening. The sequences were grouped into Operational Taxonomic Units (OTUs) that shared at least 97% sequence similarity. A representative sequence was taken for each OTU group and PyNAST (*Caporaso et al., 2010b*) was employed to align these representative sequences to the Greengenes database (*DeSantis et al., 2006*) and the taxonomic identity of each OTU was determined using the RDP Classifier (*Wang et al., 2007*). Phylotypes were considered to be contaminants if they were seen in at least two of the six negative control samples. There were 152 OTUs at the genus level present in more than one negative sample, representing 9% of the total OTUs at this level. After removing contaminant sequences and singletons, the number of quality-filtered reads per sample was between 6 and 5,861 (median = 2,306). Finally, we removed any OTUs represented by 20 or fewer reads to reduce the possibility of spurious results from the sequencing process. For among samples comparisons we rarefied each to a depth of 1,000 sequences. Thus, our assessments of microbial diversity are conservative.

Analysis of the relative diversity of microbes in extreme temperature, pH and chemical environments of homes and how it compares to habitats without each extreme condition

We compared microbial species accumulation among three extreme variables in homes: temperature, pH, and chemical extremes. Temperature was classified on a scale of 1–5, with 1 representing the coldest environments and 5 representing the hottest environments. We then binned 1 and 5 into an extreme temperature category and 2–4 into an intermediate temperature category. Similarly, environments were classified as acidic, basic or neutral and then binned into extreme pH (acidic or basic environments) *versus* neutral environments. Finally, chemical extremes were those environments characterized by the presence of detergent, bleach, metals, ammonia, or natural gas ([Table S2](#)).

We used EstimateS v.9.1.0 (*Colwell, 2013*) to construct individual-based species accumulations for all three extreme environments and their non-extreme counterparts. For these curves, reads were used as individuals and the curves were constructed using 1,000 iterations. To formally assess differences in accumulated species by read, we used $\pm 95\%$ confidence intervals for each curve. Non-overlapping 95% confidence intervals are considered formal evidence of significance (*Knezevic, 2008*).

Assessing how multiple, simultaneous extreme conditions influence microbial diversity in human homes

We were interested in testing the hypothesis that interactive effects of multiple, simultaneously extreme environmental conditions are important determinants of microbial diversity in extreme home environments (*Harrison et al., 2013*). Our study included multiple samples with more than one environmental extreme ([Table S1](#)); however, we only had sufficient replication to assess this hypothesis for 2-way interactions between extremely high temperatures and extreme pH as well as high temperature and chemical environments. Because number of reads varied significantly among different environmental extremes,

we could not use a standard 2-way ANOVA. Instead, we assessed these effects using an ordination framework.

We visualized the composition of bacteria and archaea from extreme habitats in homes using non-metric multidimensional scaling ordination (NMDS) in Primer-E v.7.0.9 with PERMANOVA + 1 (Clarke & Gorley, 2015). To do this, we first constructed NMDS plots with 100 restarts and a Type I Kruskal fit scheme based on a Dissimilarity matrix of Bray-Curtis distances. To assess the relationship between temperature (extreme vs. intermediate) and the other extremes (pH: extreme vs. neutral; chemicals: extreme vs. none) for α -diversity of microbes, we conducted a permuted multivariate analysis of variance (PERMANOVA) test with temperature class and either pH or chemical class and their interaction as factors, 9,999 iterations and Type III sums of squares. When interactions were significant (Anderson, Gorley & Clark, 2008), we conducted pairwise PERMANOVA to determine which treatment combinations significantly differed from one another. Similarly, we assessed these relationships in terms of β -diversity using a permuted dispersion (PermDisp) test of a presence/absence matrix of OTU occurrences. When these tests were significant, we conducted pairwise tests of extreme vs. non-extreme chemical and pH environments in habitats with intermediate and extreme temperatures (thus 2 tests per treatment combination). Finally, we conducted SIMPER analyses for each significant treatment combination to determine the OTUs that contributed the most to pairwise between-group differences in ordination space. Because we conducted two separate analyses for each level of diversity, we accounted for the additional error associated with multiple tests, using a revised $\alpha = 0.05/2 = 0.025$ as our cut-off for statistical significance for the results of each test. This conservative α is particularly important because we did not have equal sample sizes in all groups for these analyses, which can increase the risk of Type I error (Anderson & Walsh, 2013).

Determining which microbial genera differentiate extreme home habitats from the rest of the home

We compared the occurrences of microbes in our samples to those reported in less extreme home environments (Dunn et al., 2013). Human-associated microbes were common to both datasets, and we were particularly interested in those taxa unique to our dataset, relative to the broader home (Dunn et al., 2013). Therefore, we removed human-associated OTUs from our dataset. We identified these human-associated OTUs using databases that identified human gut (Flores et al., 2014) and skin (Urban et al., 2016) microbiomes. OTUs that occurred in at least 80% of the samples in those databases were considered human-associates and excluded from our analyses of the microbial diversity of extreme habitats in human homes. We then determined the identity of microbes that were absent from the broader homes dataset, but present in extreme environments and then tabulated the extreme habitat(s) in which they were present. Likewise, we identified the non-human associated microbes that were present in the broader home environment, but absent from all extreme environments in our samples.




Figure 1 OTU accumulation curves for each extreme environment, expressed as the number of OTUs by the number of reads from sequencing. Each curve was constructed using 1,000 iterations, and the dotted lines represent 95% confidence intervals. Non-overlapping confidence intervals indicate that the accumulation curves are significantly different. Thus, habitats with extreme temperatures had significantly more accumulated species than habitats with either extreme pH or extreme chemical environments. However, the accumulated species in habitats pH and chemical extremes did not differ significantly.

RESULTS AND DISCUSSION

What is the relative diversity of microbes in extreme temperature, pH and chemical environments of southeast US homes and how does it compare to habitats without each extreme condition?

The cumulative diversity (OTU richness) in habitats with extreme temperatures was more than twice as high as in habitats with extreme pH (maximum of 73 vs. 33, Fig. 1) and almost three times as high as habitats with extreme chemical environments (27.6; Fig. 1). Habitats with extreme temperatures also had higher OTU richness than habitats with intermediate temperatures (Fig. 2A). Conversely, previous research indicates that the diversity in habitats with either extremely high or extremely low temperatures is generally low, and dominated by a small number of abundant bacterial species (Lewin *et al.*, 2013). For example, Sharp *et al.* (2014) recently found that OTU richness in hydrothermal vents peaked at intermediate temperatures (24 °C), with reduced OTU richness in extremely hot or cold environments (Sharp *et al.*, 2014). We did not detect significant differences in the rarefied species richness of bacterial and archaeal microbes in extreme vs. neutral pH conditions; however, the marginally non-significant trend suggests that extreme pH environments also had higher microbial diversity than neutral habitats (Fig. 2B). Recent studies have demonstrated that pH is a key predictor of microbial diversity in both extreme




Figure 2 Comparison of rarefaction curves between extreme and non-extreme habitats. (A) Extreme vs. intermediate temperatures, (B) extreme vs. neutral pH environments, and (C) extreme chemicals present vs. absent. Rarefaction curves are expressed as number of OTU by number of reads from sequencing. Each curve was constructed using 1,000 iterations, and the dotted lines represent 95% confidence intervals. Significance tests were as described for Fig. 1.

environments, such as acid mine drainage sites (Kuang *et al.*, 2013), and less extreme environments, such as tropical soils (Tripathi *et al.*, 2012). In both cases, habitats with neutral pH had higher microbial diversity than those with a pH higher or lower than neutral. Thus, we again found different patterns in extreme home environments compared to other studies comparing extreme and non-extreme habitats. One possible explanation for the difference between our findings and these recent studies is that human-associated microbes are present in home environments with intermediate temperatures. Perhaps these species are able to dominate habitats with intermediate, but not extreme, conditions. Alternatively, the lower diversity in habitats with intermediate temperatures and neutral pH in our study could be due to the occurrence of extreme conditions along different axes (e.g., intermediate temperature, but extreme pH or chemical habitats). We examine potential interactive effects of these polyextreme habitats in the next section.

In contrast, habitats with extreme chemicals had significantly lower accumulated OTU richness than did habitats without these extreme conditions (Fig. 2C). Extreme chemical environments are poorly studied and understood (Rothschild & Mancinelli, 2001). Our data suggest that they could act as strong filters in extreme environments.

How do multiple, simultaneous extreme conditions influence microbial diversity in extreme home environments?

Many of the habitats in this study were characterized by more than one extreme environmental condition. Therefore, we also examined the potential for interactive effects of multiple, simultaneous extreme conditions on microbial diversity. Due to limited replication across all environmental extremes, we were only able to examine extreme pH and chemical habitats with and without extreme temperatures. We used an ordination framework to examine these interactive effects (see 'Methods').

We found significant interactions between extreme temperature and both extreme pH (PERMANOVA: Pseudo- $F_{1, 82} = 2.53$, $P = 0.0001$; Fig. 3A) and extreme chemical (PERMANOVA: Pseudo- $F_{1, 82} = 3.16$, $P = 0.0001$; Fig. 3C) environments for OTU composition. When temperatures were intermediate, there were no significant differences in microbial composition in extreme vs. neutral pH habitats (pairwise PERMANOVA: $t_{1, 38} = 1.02$, $P = 0.40$). However, when temperatures were extreme, there was a very large difference between the composition of microbes in extreme pH habitats, compared to neutral habitats (pairwise PERMANOVA: $t_{1, 38} = 1.70$, $P = 0.0001$; Fig. 3A). The five genera that contributed the most to differences between these two habitat types (from SIMPER analysis) were *Parascardovia*, *Micrococcus*, *Rothia*, *Brachybacterium*, and an unknown genus from Sphingomonadaceae. Most of these genera are associated with humans (Oshima *et al.*, 2015; Gueimonde *et al.*, 2012; Kloos & Musselwhite, 1975; Kocur, Koos & Schleifer, 2006; Vaccher *et al.*, 2007; Uchibori *et al.*, 2012). Sphingomonadaceae are widespread in aquatic habitats, including drinking water (Vaz-Moreira, Nunes & Manaia, 2011), but also other aquatic environments (e.g., tree holes-Xu *et al.*, 2008). *Brachybacterium* is usually associated with marine environments (Ward & Bora, 2006), including Antarctic sea ice (Junge *et al.*, 1998). However, it was recently detected in an urban shopping center (Tringe *et al.*, 2008). All of these genera were more common in habitats with extreme




Figure 3 NMDS ordinations OTU occurrence by (A–B) temperature & pH and (C–D) temperature & chemical environments in the home. Large symbols represent centroids ± 1 SE (A, C), and small symbols represent each sample (B, D). The interaction between temperature and pH was significant (PERMANOVA: (pseudo)- $F_{1,85} = 2.53$, $P(\text{perm}) = 0.0001$), as was the interaction between extreme temperature and chemical conditions (PERMANOVA: (pseudo)- $F_{1,85} = 3.16$, $P(\text{perm}) = 0.0001$). The ordination was constructed with Bray-Curtis distances and 100 restarts; 2-D stress was 0.21. PERMANOVA analyses were conducted using type III sums of squares and 9,999 iterations.

temperatures and neutral pH than they were in habitats with both extreme temperatures and extreme pH. While different houses had significantly different microbial composition (3-way PERMANOVA, $P = 0.0001$), there were no significant 2- or 3-way interactions with house (Table S3).

The interaction between temperature and chemical extremes was similar. Microbial composition was indistinguishable between the habitats that only had one extreme condition—regardless of whether it was temperature or chemicals that were extreme. There were also no significant differences between habitats with neither extreme temperatures nor extreme chemical conditions and habitats that had a single extreme condition. However, habitats with both extreme temperatures and extreme chemicals had significantly different microbial composition compared to all other groups (pairwise PERMANOVA; $t_{1,38} = 1.75$, $P = 0.0001$; Fig. 3C). The five genera that contributed the most to compositional difference between these two habitats (from SIMPER analysis) were *Methylobacterium*, an unknown genus of Moraxellaceae, *Sejonia*, an unknown genus of Sphingomonadaceae, and *Flavobacterium*. With the exception of the unknown genus of Moraxellaceae, which was more common in extreme chemical and temperature environments, all of these genera were more common in the habitats without temperature and chemical extremes. Moraxellaceae have been found in other extreme environments, including deep sea sediments




Figure 4 Average distances between samples and centroids (β -diversity) across home environments that differ with respect to extreme temperatures and (A) extreme pH conditions & (B) extreme chemical conditions. Data were assessed using PermDisp; dispersion was significantly different across extreme temperatures and extreme pH conditions ($F_{3,82} = 4.08, P = 0.024$) and across extreme temperatures and extreme chemical conditions ($F_{3,82} = 6.99, P = 0.0017$). Post-hoc pairwise tests: * $P < 0.025$, ** $P < 0.01$, *** $P < 0.001$.

(Maruyama et al., 1997). Although it was more common in our less extreme environments, *Sejonia* is better known from Antarctic ice (Yi, Yoon & Chun, 2005). Sphingomonadaceae as described above are common to aquatic habitats. *Methylobacterium* is a widespread habitat generalist that is facultatively methyltrophic (Green, 2006). Finally, *Flavobacterium* is common in freshwater and marine ecosystems but tends to flourish in cold environments with high salinity (Bernardet & Bowman, 2006).

There were also significant differences in the β -diversity in home environments with more than one extreme condition. When temperatures were intermediate, there were no significant differences between neutral and extreme pH environments (Fig. 4A; PermDisp: $P = 0.3864$). However when temperatures were also extreme, habitats with extreme pH conditions had significantly higher β -diversity than those with neutral pH conditions (Fig. 4A; PermDisp: $P = 0.0014$). Similarly, at intermediate temperatures, there was a non-significant trend (Fig. 4B; PermDisp: $P = 0.03$, Bonferroni-corrected $\alpha = 0.025$) in which habitats without extreme chemicals present had higher β -diversity than those with extreme chemicals present. However, when temperatures were also extreme, habitats with extreme chemicals present had higher β -diversity than those without extreme chemicals (Fig. 4B; PermDisp: $P = 0.0006$). This increase in β -diversity in extreme pH and chemical environments when temperatures were also extreme suggests that polyextreme conditions may support a higher diversity of extremophiles and/or reduced occurrences of numerically dominant genera compared to environments with a single extreme condition, at least among habitats (in contrast to within habitats). The 5 genera that contributed the most to differences in β -diversity between neutral and extreme pH conditions when temperatures were also extreme were: *Veillonella*, *Kocuria*, *Peptoniphilus*, *Parascardovia*,

Table 1 Summary of occurrences of microbes that were present in samples from extreme home environments, but absent from the broader home samples. Numbers indicate the number of reads of each genus by extreme environment. The first group includes genera that were only present in one extreme environment, the second group includes genera that were present in two extreme environments, and the last group includes genera that were present in all three extreme home environments.

Genus	Extreme temperatures	Extreme pH	Extreme Chemical
<i>Brochothrix</i>	265	0	0
<i>Buchnera</i>	22	0	0
<i>Polynucleobacter</i>	33	0	0
<i>Ralstonia</i>	21	0	0
<i>Thermicanus</i>	34	0	0
<i>Helcococcus</i>	0	0	22
<i>Solibacter</i>	86	0	30
<i>Brevundimonas</i>	184	189	0
<i>Azobacteroides</i>	0	33	33
<i>Elizabethkingia</i>	0	25	24
<i>Xiphinematobacter</i>	0	19	21
<i>Azospira</i>	139	33	44
<i>Brachybacterium</i>	101	52	69
<i>Enhydrobacter</i>	452	387	408
<i>Gluconobacter</i>	23	21	22
<i>Oligella</i>	40	74	77
<i>Parascardovia</i>	141	46	107
<i>Photobacterium</i>	71	65	93
<i>Propionibacterium</i>	73	31	40
<i>Salinibacterium</i>	108	334	355

and *Anaerococcus*. Interestingly, these were also the top 5 genera contributing to differences between habitats with and without extreme chemicals that also had extreme temperatures. All of these genera were less common in habitats with 2 extremes than they were in habitats with only extreme temperatures. They are also genera that include human-associated species (Bhatti & Frank, 2000; Fadda, Vignolo & Oliver, 2001; Song, Liu & Finegold, 2007; Gueimonde et al., 2012).

Which microbial genera differentiate extreme home habitats from the rest of the home?

After removing all human-associated microbes (see 'Methods'), there were a total of 241 unique genera in the broader homes dataset (Dunn et al., 2013). Our extreme samples contained 135 of the remaining broader homes genera, but ~44% of the genera found in the broader homes were absent from our extreme home samples (Table S4), the absence of which might simply be due to the larger number of samples in Dunn et al. (2013). More interestingly, we found 20 genera present among our samples that were absent from the broader homes dataset. Nine of these genera were found in all three categories of extreme environments (Table 1); one genus (*Solibacter*) was absent from habitats with extreme pH, but occurred in both extreme chemical and temperature environments. *Solibacter* is

a common and abundant soil microbe, especially in tropical regions ([Guan et al., 2013](#); [Wang et al., 2015](#)). There was also one genus (*Brevundimonas*) that was absent from extreme chemical environments, but present in both extreme temperature and extreme pH environments; *Brevundimonas* is one of the only genera thought to be able to survive the low temperatures and ionizing radiation on Mars ([Dartnell et al., 2010](#)). There were three genera (*Azobacteroides*, *Elizabethkingia*, and *Xiphinematobacter*) that occurred in both extreme pH and chemical environments that were absent in extreme temperature environments. Both *Azobacteroides* and *Xiphinematobacter* are gut symbionts of invertebrates; *Azobacteroides* is commonly found inside the protozoan symbionts of termites ([Noda et al., 2007](#)), and *Xiphinematobacter* is an endosymbiont of nematodes ([Vandekerckhove et al., 2000](#)). In invertebrate guts these microbes likely experience extreme chemical and pH environments frequently, while being relatively protected from temperature stress. *Elizabethkingia* is a cosmopolitan genus, with species that are endosymbionts of mosquitoes ([Kämpfer et al., 2011](#)), and others that are pathogens of both humans ([Ceyhan & Celik, 2011](#)) and frogs ([Xie et al., 2009](#)). There was one genus that was only found in extreme chemical environments (*Helcococcus*). Interestingly, members of the genus *Helcococcus* possess the ability to degrade detergents. In fact, the detergent Tween-80 can be added to media to enrich *Helcococcus* ([Collins et al., 1993](#); [Chagla et al., 1998](#)). Finally, we found 5 genera (*Brochothrix*, *Buchnera*, *Polynucleobacter*, *Ralstonia*, and *Thermicanus*) unique to extreme temperature environments. *Brochothrix* is a common spoilage bacterium in meat ([Rattanasomboon et al., 1999](#)). *Buchnera* is a widespread aphid endosymbiont ([Shigenobu et al., 2000](#)). Recently, a survey of homes in Raleigh, NC demonstrated that aphids could be quite common in human homes ([Bertone et al., 2016](#)), which could explain how this genus arrived in the homes in our study (via aphids in the home). The genus *Polynucleobacter* includes both free-living species and species that are endosymbionts of nematodes ([Vannini et al., 2007](#)). *Ralstonia metallidurans* is a bacterium specifically adapted to toxic metal environments ([Mergeay et al., 2003](#)). Other species of *Ralstonia* have been shown to be effectively controlled using high temperature treatments in commercial crops ([Kongkiattikajorn & Thepa, 2007](#)). In our study, *Ralstonia* were collected in both high and low temperature environments. Finally, *Thermicanus* is, as its name suggests, a thermophilic bacterial genus ([Wrighton et al., 2008](#)).

CONCLUSIONS

This study has provided a glimpse into the microbial diversity that lives in habitats of human homes similar in their extreme temperature, pH and chemical conditions to some of the most extreme habitats on Earth. We discovered that these conditions have lower diversity than the surrounding home environment; yet tens of bacterial lineages can be found in these extreme habitats of the human home, including many taxa with known associations with extreme conditions. Habitats with extreme temperatures alone appear to be able to support a greater diversity of microbes than habitats with extreme pH or extreme chemical environments alone. Microbial diversity is significantly lowest when habitats have both extreme temperature and one of these other extremes. Interestingly, environments

in homes often alternate between periods of extreme and non-extreme conditions. For example, dishwashers are only likely to have extremely high temperatures while cleaning and drying dishes. This variability could lead to temporal shifts in microbial composition, similar to those found for human vaginal microbes (*Gajer et al., 2012*). This variability may also explain the presence of human-associated generalist species in our samples. Future work, with samples taken before and after appliances (like many of those used in our study) are operated, could elucidate the importance of episodic extreme conditions for microbial communities in homes. Additionally, a key next step is understanding which of the relatively few species that are found in these poly-extreme environments in the home are metabolically active there and both whether these polyextreme taxa pose health threats (as was recently suggested by *Gümral et al., 2016*) and/or might be useful industrially.

ACKNOWLEDGEMENTS

We thank the homeowners for allowing us to sample habitats in their homes. Thanks also to Dr. Holly Menninger for her support and guidance in addressing logistical challenges presented by the current work. The Genomics Laboratory at the North Carolina Museum of Natural Sciences provided critical logistical support for this project.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was funded by A.P. Sloan Microbiology of the Built Environment Program grant awarded to RRD. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
A.P. Sloan Microbiology of the Built Environment Program.

Competing Interests

The authors do not have any competing financial or non-financial competing interests to report.

Author Contributions

- Amy M. Savage analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables.
- Justin Hills and Katherine Driscoll conceived and designed the experiments, performed the experiments, reviewed drafts of the paper.
- Daniel J. Fergus and Amy M. Grunden contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Robert R. Dunn conceived and designed the experiments, contributed reagents/materials/analysis tools, reviewed drafts of the paper.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:
NCBI [SRP071677](#).

Data Availability

The following information was supplied regarding data availability:
The raw data has been supplied as a [Supplemental Dataset](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2376#supplemental-information>.

REFERENCES

- Anderson MJ, Gorley RN, Clark KR. 2008. *PERMANOVA + for PRIMER: guide to software and statistical methods*. Plymouth: PRIMER-E.
- Anderson MJ, Walsh DCI. 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of heterogeneous dispersions: what null hypothesis are you testing? *Ecological Monographs* 23:557–574.
- Bates ST, Berg-Lyons D, Caporaso JG, Walters WA, Knight R, Fierer N. 2011. Examining the global distribution of dominant archaeal populations in soil. *The ISME Journal* 5:908–917 DOI [10.1038/ismej.2010.171](https://doi.org/10.1038/ismej.2010.171).
- Bernardet JF, Bowman JP. 2006. The genus *Flavobacterium*. In: *The prokaryotes*. New York: Springer, 481–531.
- Bertone MA, Leong M, Bayless KM, Malow TLF, Dunn RR, Trautwein MD. 2016. Arthropods of the great indoors: characterizing diversity inside urban and suburban homes. *PeerJ* 4:e1582 DOI [10.7717/peerj.1582](https://doi.org/10.7717/peerj.1582).
- Bhatti MA, Frank MO. 2000. *Veillonella parvula* meningitis: case report and review of *Veillonella* infections. *Clinical Infectious Diseases* 3:839–840.
- Boe-Hansen R, Albechtsen H-J, Arvin E, Jorgensen C. 2002. Bulk water phase and biofilm growth in drinking water at low nutrient conditions. *Water Research* 36:4477–4486 DOI [10.1016/S0043-1354\(02\)00191-4](https://doi.org/10.1016/S0043-1354(02)00191-4).
- Brock TD, Boylen KL. 1973. Presence of thermophilic bacteria in laundry and domestic hot water heaters. *Applied Microbiology* 25:72–76.
- Caporaso JG, Bittinger K, Bushman FD, Desantis TZ, Andersen GL, Knight R. 2010b. PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* 26:266–267 DOI [10.1093/bioinformatics/btp636](https://doi.org/10.1093/bioinformatics/btp636).
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon J, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunencko T, Zaneveld J, Knight R. 2010a. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7:335–336 DOI [10.1038/nmeth.f.303](https://doi.org/10.1038/nmeth.f.303).

- Ceyhan M, Celik M. 2011. *Elizabethkingia meningosepticum* (*Chryseobacterium meningosepticum*) infections in children. *International Journal of Pediatrics* 2011: Article 215237 DOI 10.1155/2011/215237.
- Chagla AH, Borczyk AA, Facklam RR, Lovgren M. 1998. Breast abscess associated with *Helcococcus kunzii*. *Journal of Clinical Microbiology* 36:2377–2379.
- Clarke KR, Gorley RN. 2015. Primer v.7: user manual/tutorial. Plymouth: Primver-E.
- Collins MD, Facklam RR, Rodrigues UM, Ruoff KL. 1993. Phylogenetic analysis of some *Aerococcus*-like organisms from clinical sources: description of *Helcococcus kunzii* gen. nov., sp. nov. *International Journal of Systematic Bacteriology* 43:425–429 DOI 10.1099/00207713-43-3-425.
- Colwell RK. 2013. *EstimateS*: statistical estimation of species richness and shared species from samples. Version 9. Available at purl.oclc.org/estimates.
- Dartnell LR, Hunter SJ, Lovell KV, Coates AJ, Ward JM. 2010. Low temperature ionizing radiation resistance of *Deinococcus radiodurans* and Antarctic Dray Valley bacteria. *Astrobiology* 7:717–732 DOI 10.1089/ast.2009.0439.
- DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen GL. 2006. Greengenes, a Chimera-Checked 16S rRNA gene database and workbench compatible with ARB. *Applied Environmental Microbiology* 72:5069–5072 DOI 10.1128/AEM.03006-05.
- Dunn RR, Fierer N, Henley JB, Leff JW, Menninger HL. 2013. Home life: factors structuring the bacterial diversity found within and between homes. *PLoS ONE* 8(5):e64133 DOI 10.1371/journal.pone.0064133.
- Elleuche S, Schröder C, Sahn K, Antranikian G. 2014. Extremozymes-biocatalysts with unique properties from extremophilic microorganisms. *Current Opinion in Biotechnology* 29:116–123 DOI 10.1016/j.copbio.2014.04.003.
- Fadda S, Vignolo G, Oliver G. 2001. Tyramine degradation and tyramine/histamine production by lactic acid bacteria and *Kocuria* strains. *Biotechnology Letters* 23:2015–2019.
- Fierer N, Hamady M, Lauber CL, Knight R. 2008. The influence of sex, handedness, and washing on the diversity of hand surface bacteria. *Proceedings of the National Academy of Sciences of the United States of America* 105:17994–17999 DOI 10.1073/pnas.0807920105.
- Flores GE, Caporaso JG, Henley JB, Rideout JR, Domogala D, Chase J, Leff JW, Vásquez-Baeza Y, Gonzalez A, Knight R, Dunn RR, Fierer N. 2014. Temporal variability is a personalized feature of the human microbiome. *Genome Biology* 15: 531 DOI 10.1186/s13059-014-0531-y.
- Gajer P, Brotman RM, Bai G, Sakamoto J, Schütte UME, Zhong X, Koenig SK, Fu L, Ma Z, Zhou X, Abdo Z, Forney LJ, Ravel J. 2012. Temporal dynamics of the human vaginal microbiota. *Science Translational Medicine* 4(132):132ra52 DOI 10.1126/scitranslmed.3003605.
- Green PN. 2006. *Methylobacterium*. In: *The prokaryotes*. New York: Springer, 257–265.

- Guan X, Wang J, Zhao H, Wang J, Luo X, Zhao F. 2013.** Soil bacterial communities shaped by geochemical factors and land use in a less-explored area, Tibetan Plateau. *BMC Genomics* **14**:820 DOI [10.1186/1471-2164-14-820](https://doi.org/10.1186/1471-2164-14-820).
- Gueimonde M, Bottacini F, Van Sinderen D, Ventura M, Margolles A, Sanchez B. 2012.** Genome sequence of *Parascardovia denticolens* IPLA 20019, isolated from human breast milk. *Journal of Bacteriology* **194**:4776–4777 DOI [10.1128/JB.01035-12](https://doi.org/10.1128/JB.01035-12).
- Gümral R, Özhak-Baysan B, Tümgör A, Saraçlı MA, Yidrian ŞT, Ilkit M, Zupančič J, Novak-Babič M, Gunde-Cimerman N, Zalar P, De Hoog GS. 2016.** Dishwashers provide a selective extreme environment for human-opportunistic yeast-like fungi. *Fungal Diversity* **76**:1–9 DOI [10.1007/s13225-015-0327-8](https://doi.org/10.1007/s13225-015-0327-8).
- Harrison JP, Gheeraert N, Tsigelnitskiy D, Cockell CS. 2013.** The limits of life under multiple extremes. *Trends in Microbiology* **21**:204–212 DOI [10.1016/j.tim.2013.01.006](https://doi.org/10.1016/j.tim.2013.01.006).
- Hoehler TM, Jorgensen BB. 2013.** Microbial life under extreme energy limitation. *Nature Reviews Microbiology* **11**:83–95 DOI [10.1038/nrmicro2939](https://doi.org/10.1038/nrmicro2939).
- Hulcr J, Latimer AM, Henley JB, Rountree NR, Fierer N, Lucky A, Lowman MD, Dunn RR. 2012.** A jungle in there: bacteria in belly buttons are highly diverse, but predictable. *PLoS ONE* **7**(11):e47712 DOI [10.1371/journal.pone.0047712](https://doi.org/10.1371/journal.pone.0047712).
- Junge K, Gosink JJ, Hoppe HG, Staley JT. 1998.** *Arthrobacter*, *Brachybacterium* and *Planococcus* isolates identified from Antarctic sea ice brine. *Systematic and Applied Microbiology* **21**:306–314 DOI [10.1016/S0723-2020\(98\)80038-6](https://doi.org/10.1016/S0723-2020(98)80038-6).
- Kalmbach S, Manz W, Szewzyk U. 1997.** Isolation of new bacterial species from drinking water biofilms and proof of their *in situ* dominance with highly specific 16S rRNA probes. *Applied and Environmental Microbiology* **63**:4164–4170.
- Kämpfer P, Matthews H, Glaeser SP, Martin K, Lidders N, Faye I. 2011.** *Elizabethkingia anophelis* sp. nov., isolated from the midgut of the mosquito *Anopheles gambiae*. *International Journal of Systematic and Evolutionary Microbiology* **61**:2670–2675 DOI [10.1099/ijs.0.026393-0](https://doi.org/10.1099/ijs.0.026393-0).
- Kloos WE, Musselwhite MS. 1975.** Distribution and persistence of *Staphylococcus* and *Micrococcus* species and other aerobic bacteria on human skin. *Applied Microbiology* **30**:381–395.
- Knezevic A. 2008.** *Overlapping confidence intervals and statistical significance*. StatNews: Cornell University Statistical Consulting Unit 73.
- Kocur M, Koos WE, Schleifer KH. 2006.** The genus *Micrococcus*. *Prokaryotes* **3**:961–971.
- Kongkiattikajorn J, Thepa S. 2007.** Increased tomato yields by heat treatment for controlling *Ralstonia solanacearum* in soil. In: *Proceedings of the 45th Kasetsart University Annual Conference*.
- Kuang J-L, Huang L-N, Chen L-X, Hua Z-S, Li S-J, Shu W-S. 2013.** Contemporary environmental variation determines microbial diversity patterns in acid mine drainage. *International Society for Microbial Ecology* **7**:1038–1050 DOI [10.1038/ismej.2012.139](https://doi.org/10.1038/ismej.2012.139).
- Lauber CL, Hamady M, Knight R, Fierer N. 2009.** Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applied and Environmental Microbiology* **75**:5111–5120 DOI [10.1128/AEM.00335-09](https://doi.org/10.1128/AEM.00335-09).

- Lewin A, Wentzel A, Valla S. 2013. Metagenomics of microbial life in extreme temperature environments. *Current Opinion in Biotechnology* 24:516–525 DOI 10.1016/j.copbio.2012.10.012.
- Martin LJ, Adams RI, Bateman A, Bik HM, Hawks J, Hird SM, Hughes D, Kembel SW, Kinney K, Kolokotronis SO, Levy G, McClain C, Meadow JF, Medina RF, Mhuireach G, Moreau CS, Munshi-South J, Nichols LM, Palmer C, Popova L, Schal C, Taubel M, Trautwein M, Ugalde JA, Dunn RR. 2015. Evolution of the indoor biome. *Trends in Ecology and Evolution* 1909:1–10 DOI 10.1016/j.tree.2015.02.001.
- Maruyama A, Taniguchi R, Tanaka H, Ishiwata H, Higashihara T. 1997. Low-temperature adaptation of deep-sea bacteria isolated from the Japan Trench. *Marine Biology* 128:705–711 DOI 10.1007/s002270050138.
- Mergeay M, Monchy S, Vallaeyts T, Auquier V, Benotmane A, Bertin P, Taghavi S, Dunn J, Van der Lelie D, Wattiez R. 2003. *Ralstonia metallidurans*, a bacterium specifically adapted to toxic metals: towards a catalogue of metal responsive genes. *FEMS Microbiology Reviews* 27:385–410 DOI 10.1016/S0168-6445(03)00045-7.
- Niehaus F, Bertolodo C, Kähler M, Antranikian G. 1999. Extremophiles as a source of novel enzymes for industrial application. *Applied Microbiology and Biotechnology* 51:711–729 DOI 10.1007/s002530051456.
- Noda S, Kitade O, Inoue T, Kawi M, Kanuka M, Hiroshima K, Hongoh Y, Constantino R, Uys V, Zhong J, Kudo T, Ohkuma M. 2007. Cospeciation in the triplex symbiosis of termite gut protists (*Pseudotrichonympha* spp.), their hosts, and their bacterial endosymbionts. *Molecular Ecology* 16:1257–1266 DOI 10.1111/j.1365-294X.2006.03219.x.
- Oshima K, Hayashi J, Toh H, Nakano A, Shindo C, Komiya K, Morita H, Honda K, Hattori M. 2015. Complete genome sequence of *Parascardovia denticolens* JCM 12538^T, isolated from human dental caries. *Genome Announcements* 3(3):e00485–15.
- Rattanasomboon N, Bellara SR, Harding CL, Fryer PJ, Thomas CR, Al-Rubeai M, McFarlane CM. 1999. Growth and enumeration of the meat spoilage bacterium *Brochothrix thermosphacta*. *International Journal of Food Microbiology* 51:145–158 DOI 10.1016/S0168-1605(99)00129-4.
- Rothschild LJ, Mancinelli RL. 2001. Life in extreme environments. *Nature* 409:1092–1101 DOI 10.1038/35059215.
- Sharp CE, Brady AL, Sharp GH, Grasby SE, Stott MB, Dunfield PF. 2014. Humboldt's spa: microbial diversity is controlled by temperature in geothermal environments. *ISME Journal* 8:1166–1174 DOI 10.1038/ismej.2013.237.
- Shigenobu S, Watanabe H, Hattori M, Sakaki Y, Ishikawa H. 2000. Genome sequence of the endocellular bacterial symbiont of aphids Buchnera sp. APS. *Nature* 407:81–86.
- Song Y, Liu C, Finegold SM. 2007. *Peptoniphilus gorbachii* sp. nov., *Peptoniphilus olsenii* sp. nov., and *Anaerococcus murdochii* sp. nov. isolated from clinical specimens of human origin. *Journal of Clinical Microbiology* 45:1746–1752.
- Szewzyk U, Szewzyk R, Manz W, Schleifer KH. 2000. Microbiological safety of drinking water. *Annual Review of Microbiology* 54:81–127 DOI 10.1146/annurev.micro.54.1.81.

- Tringe SF, Zhang T, Liu X, Yu Y, Lee WH, Yap J, Yao F, Suan ST, Ing SK, Haynes M, Rohwer F, Wei CL, Tan P, Bristow J, Rubin EM, Ruan Y. 2008. The air-borne metagenome in an indoor urban environment. *PLoS ONE* 3(4):e1862 DOI 10.1371/journal.pone.0001862.
- Tripathi BM, Kim M, Singh D, Lee-Cruz L, Lai-Hoe A, Ainuddin AN, Go R, Rahim RA, Husni MHA, Chun J, Adams JM. 2012. Tropical soil bacterial communities in Malaysia: pH dominates in the equatorial tropics too. *Microbial Ecology* 64:474–484 DOI 10.1007/s00248-012-0028-8.
- Uchibori S, Tsuzukibashi O, Kobayashi T, Aida M. 2012. Localization of the genus *Rothia* in the oral cavity. *International Journal of Oral-Medical Science* 11:207–210 DOI 10.5466/ijoms.11.207.
- Urban JM, Fergus DJ, Savage AM, Ehlers M, Menninger HL, Dunn RR, Horvath JE. 2016. The effect of habitual antiperspirant and deodorant product use on the armpit microbiome. *PeerJ* 4:e1605 DOI 10.7717/peerj.1605.
- Vaccher S, Cordiali R, Osimani P, Manso E, De Benedictis FM. 2007. Bacteremia caused by *Rothia mucilaginosa* in a patient with Shwachman-Diamond Syndrome. *Infection* 35:209 DOI 10.1007/s15010-007-6284-8.
- Valentine DL. 2007. Adaptations to energy stress dictate the ecology and evolution of Archea. *Nature Reviews Microbiology* 5:316–323 DOI 10.1038/nrmicro1619.
- Vandekerckhove TTM, Willems A, Gillis M, Coomans A. 2000. Occurrence of novel verrucomicrobial species endosymbiotic and associated with parthenogenesis in *Xiphinema americanum*-group species (Nematoda, Longidoridae). *International Journal of Systematic and Evolutionary Microbiology* 50:2197–2205 DOI 10.1099/00207713-50-6-2197.
- Van den Burg B. 2003. Extremophiles as a source of novel enzymes. *Current Opinion in Microbiology* 6:213–218 DOI 10.1016/S1369-5274(03)00060-2.
- Vannini C, Pöckl M, Petroni G, Wu QL, Lang E, Stackebrandt E, Schrollhammer M, Richardson PM, Hahn MW. 2007. Endosymbiosis in *Statu nascendi*: close phylogenetic relationship between obligately endosymbiotic and obligately free-living *Polynucleobacter* strains (Betaproteobacteria). *Environmental Microbiology* 9:347–359 DOI 10.1111/j.1462-2920.2006.01144.x.
- Vaz-Moreira I, Nunes OC, Manaia CM. 2011. Diversity and antibiotic resistance patterns of *Sphingomonadaceae* isolates from drinking water. *Applied and Environmental Microbiology* 77:5697–5706 DOI 10.1128/AEM.00579-11.
- Wang N, Ding LJ, Xu HJ, Li HB, Su JQ, Zhu YG. 2015. Variability in responses of bacterial communities and nitrogen oxide emission to urea fertilization among various flooded paddy soils. *FEMS Microbiology Ecology* 91:fiv013 DOI 10.1093/femsec/fiv013.
- Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naïve Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied And Environmental Microbiology* 73:5261–5267 DOI 10.1128/AEM.00062-07.
- Ward AC, Bora N. 2006. Diversity and biogeography of marine actinobacteria. *Current Opinion in Microbiology* 9:279–286.

- Wrighton KC, Agbo P, Warnecke F, Weber KA, Brodie EL, DeSantis TZ, Hugenholtz P, Andersen GL, Coates JD. 2008.** A novel ecological role of the Firmicutes identified in thermophilic microbial fuel cells. *ISME Journal* 2:1146–1156 DOI [10.1038/ismej.2008.48](https://doi.org/10.1038/ismej.2008.48).
- Xie ZY, Zhou YC, Wang SF, Mei B, Xu XD, Wen WY, Feng YQ. 2009.** First isolation and identification of *Elizabethkingia meningoseptica* from cultured tiger frog, *Rana tigrina rugulosa*. *Veterinary Microbiology* 138:140–144.
- Xu Y, Chen S, Kaufman MG, Maknojia S, Bagdasarian M, Walker ED. 2008.** Bacterial community structure in tree hole habitats of *Ochlerotatus triseriatus*: influences of larval feeding. *Journal of the American Mosquito Association* 24:219–227.
- Yi H, Yoon HI, Chun J. 2005.** *Sejongia antarctica* gen. nov., sp. nov. and *Sejongia jeonii* sp. nov., isolated from the Antarctic. *International Journal of Systematic and Evolutionary Microbiology* 55:409–416.